

Physics 111 - Class 5A

Forces I


October 3, 2022

Class Outline

- Logistics / Announcements
- Introduction to Chapter 5
- Clicker Questions
- Activity: Worked Problem

Logistics/Announcements

- Lab this week
- HW5 due this week on Thursday at 6 PM
- Learning Log 5 due on Saturday at 6 PM
- HW and LL deadlines have a 48 hour grace period
- Test/Bonus Test: Test 2 available this week (Chapters 3 & 4)
 - Test will be on Friday during class! You must be physically present to write it.



Physics 111

Search this book...

Unsyllabus

ABOUT THIS COURSE

[Course Syllabus \(Official\)](#)

[Course Schedule](#)

[Accommodations](#)

[How to do well in this course](#)

GETTING STARTED

[Before the Term starts](#)

[After the first class](#)

[In the first week](#)

[Week 1 - Introductions!](#)

PART 1 - KINEMATICS

[Week 2 - Chapter 2](#)

[Week 3 - Chapter 3](#)

[Week 4 - Chapter 4](#)

PART 2 - DYNAMICS

[Week 5 - Chapter 5](#)

[Readings](#)

[Videos](#)


[Homework](#)

[Week 5 Classes](#)

[Test](#)

Content Summary from Crash Course Physics

Newton's Laws



Required Videos

1. Introduction to Inertia and Inertial Mass



Checklist of items

- ☐ Video 1
- ☐ Video 2
- ☐ Video 3
- ☐ Video 4
- ☐ Video 5
- ☐ Video 6
- ☐ Video 7
- ☐ Video 8
- ☐ Video 9
- ☐ Video 10
- ☐ Video 11
- ☐ Video 12

Introduction

☰ Table of contents



Preface

▼ Mechanics

- ▶ 1 Units and Measurement
- ▶ 2 Vectors
- ▶ 3 Motion Along a Straight Line
- ▶ 4 Motion in Two and Three Dimensions
- ▼ 5 Newton's Laws of Motion

Introduction

5.1 Forces

5.2 Newton's First Law

5.3 Newton's Second Law

5.4 Mass and Weight

5.5 Newton's Third Law

5.6 Common Forces

5.7 Drawing Free-Body Diagrams

▶ Chapter Review

- ▶ 6 Applications of Newton's Laws
- ▶ 7 Work and Kinetic Energy
- ▶ 8 Potential Energy and Conservation of Energy
- ▶ 9 Linear Momentum and Collisions
- ▶ 10 Fixed-Axis Rotation
- ▶ 11 Angular Momentum

Search this book



📖 My highlights



Figure 5.1 The Golden Gate Bridge, one of the greatest works of modern engineering, was the longest suspension bridge in the world in the year it opened, 1937. It is still among the 10 longest suspension bridges as of this writing. In designing and building a bridge, what physics must we consider? What forces act on the bridge? What forces keep the bridge from falling? How do the towers, cables, and ground interact to maintain stability?

Chapter Outline

[5.1 Forces](#)

[5.2 Newton's First Law](#)

[5.3 Newton's Second Law](#)

[5.4 Mass and Weight](#)

[5.5 Newton's Third Law](#)

[5.6 Common Forces](#)

[5.7 Drawing Free-Body Diagrams](#)

“Kinematics” and “Dynamics”

- For the past two weeks we have been talking about “Kinematics” in 1D, 2D, and 3D while we studied vectors...
- Kinematics help us describe the way objects move
 - You were told objects had this initial velocity, or that acceleration etc...
 - But how did they get those initial velocities ?
- The next two chapters are about “Dynamics” - how forces affect the motion of objects and systems.

Monday's Class

5.1 Forces

5.2 Newton's First Law

5.4 Mass and Weight

5.7 Free Body Diagrams

Units of Force

The equation $F_{\text{net}} = ma$ is used to define net force in terms of mass, length, and time. As explained earlier, the SI unit of force is the newton. Since $F_{\text{net}} = ma$,

$$1 \text{ N} = 1 \text{ kg} \cdot \text{m/s}^2.$$

Although almost the entire world uses the newton for the unit of force, in the United States, the most familiar unit of force is the pound (lb), where $1 \text{ N} = 0.225 \text{ lb}$. Thus, a 225-lb person weighs 1000 N.

WEIGHT

The gravitational force on a mass is its weight. We can write this in vector form, where $\vec{w}$ is weight and m is mass, as

$$\vec{w} = m\vec{g}. \quad 5.8$$

In scalar form, we can write

$$w = mg. \quad 5.9$$

Adding Forces together

Free Body Diagram




Figure 5.3

(a) An overhead view of two ice skaters pushing on a third skater. Forces are vectors and add like other vectors, so the total force on the third skater is in the direction shown.

Drawing Free Body Diagrams

Drawing Free-Body Diagrams


1. Draw the object under consideration. If you are treating the object as a particle, represent the object as a point. Place this point at the origin of an xy -coordinate system.
2. Include all forces that act on the object, representing these forces as vectors. However, do not include the net force on the object or the forces that the object exerts on its environment.
3. Resolve all force vectors into x - and y -components.
4. Draw a separate free-body diagram for each object in the problem.

Drawing Free Body Diagrams


Drawing Free-Body Diagrams

1. Draw the object under consideration. If you are treating the object as a particle, represent the object as a point. Place this point at the origin of an xy -coordinate system.
2. Include all forces that act on the object, representing these forces as vectors. However, do not include the net force on the object or the forces that the object exerts on its environment.
3. Resolve all force vectors into x - and y -components.
4. Draw a separate free-body diagram for each object in the problem.

Drawing Free Body Diagrams



(a) Box at rest on a horizontal surface



(b) Box on an inclined plane

Figure 5.4 In these free-body diagrams, $\vec{N}$ is the normal force, $\vec{W}$ is the weight of the object, and $\vec{f}$ is the friction.

Sled Pulled at an Angle




Figure 5.31

A sled is pulled by force P at an angle of 30° . Draw a Free Body Diagram of all the forces on the sled, and resolve the forces into their x and y components.

Sled Pulled at an Angle

Free Body Diagram




Figure 5.31

A sled is pulled by force P at an angle of 30° . Draw a Free Body Diagram of all the forces on the sled, and resolve the forces into their x and y components.

Sled Pulled at an Angle




Figure 5.31

A sled is pulled by force P at an angle of 30° . Draw a Free Body Diagram of all the forces on the sled, and resolve the forces into their x and y components.

Blocks on a Ramp






Figure 5.32

Block A is resting on top of Block B and the two blocks are placed on a ramp with an angle θ . Draw the Free Body Diagrams of both blocks, and resolve the components.

Blocks on a Ramp

Free Body Diagram



Free Body Diagram





Figure 5.32

Block A is resting on top of Block B and the two blocks are placed on a ramp with an angle θ . Draw the Free Body Diagrams of both blocks, and resolve the components.

Two Blocks in Contact




Free Body Diagram

Example 5.15


Block 1 (mass m_1) is in contact with Block 2 (mass m_2) and a force F is applied to m_1 , towards the right.

Draw the Free Body Diagrams on both masses.

Two Blocks in Contact



Free Body Diagram



Example 5.15

Block 1 (mass m_1) is in contact with Block 2 (mass m_2) and a force F is applied to m_1 , towards the right.

Draw the Free Body Diagrams on both masses.

Key Equations

Net external force	$\vec{\mathbf{F}}_{\text{net}} = \sum \vec{\mathbf{F}} = \vec{\mathbf{F}}_1 + \vec{\mathbf{F}}_2 + \dots$
Newton's first law	$\vec{\mathbf{v}} = \text{constant when } \vec{\mathbf{F}}_{\text{net}} = \vec{\mathbf{0}} \text{ N}$
Newton's second law, vector form	$\vec{\mathbf{F}}_{\text{net}} = \sum \vec{\mathbf{F}} = m\vec{\mathbf{a}}$
Newton's second law, scalar form	$F_{\text{net}} = ma$
Newton's second law, component form	$\sum \vec{\mathbf{F}}_x = m\vec{\mathbf{a}}_x, \sum \vec{\mathbf{F}}_y = m\vec{\mathbf{a}}_y, \text{ and } \sum \vec{\mathbf{F}}_z = m\vec{\mathbf{a}}_z.$
Newton's second law, momentum form	$\vec{\mathbf{F}}_{\text{net}} = \frac{d\vec{\mathbf{p}}}{dt}$
Definition of weight, vector form	$\vec{\mathbf{w}} = m\vec{\mathbf{g}}$
Definition of weight, scalar form	$w = mg$

Key Equations

Newton's third law	$\vec{\mathbf{F}}_{\text{AB}} = -\vec{\mathbf{F}}_{\text{BA}}$
Normal force on an object resting on a horizontal surface, vector form	$\vec{\mathbf{N}} = -m\vec{\mathbf{g}}$
Normal force on an object resting on a horizontal surface, scalar form	$N = mg$
Normal force on an object resting on an inclined plane, scalar form	$N = mg\cos\theta$
Tension in a cable supporting an object of mass m at rest, scalar form	$T = w = mg$

Clicker Questions

CQ.5.1

An object weighs 294 N on Earth. What is its weight on the Moon?

a) 50.1 N

b) 30.0 N

c) 249 N

d) 1461 N

$$g_{moon} \approx \frac{g_{earth}}{6}$$

A

B

C

D

E

CQ.5.2

Two people apply the same force from the same height to throw two identical balls in the air. Will the balls necessarily travel the same distance? Why or why not?

- a) No, the balls will not necessarily travel the same distance because the gravitational force acting on them is different.
- b) No, the balls will not necessarily travel the same distance because the angle at which they are thrown may differ.
- c) Yes, the balls will travel the same distance because the gravitational force acting on them is the same.
- d) Yes, the balls will travel the same distance because the angle at which they are thrown may differ.

A

B


C

D

E

CQ.5.3

An object slides down an inclined plane as in this figure. A frictional force acts on the block. Which of these statements is true?



- I. The frictional force is up the inclined plane.
- II. The frictional force depends on the mass of the object.
- III. If θ increases, the frictional force will always decrease.
- IV. The frictional force is dependent on the speed of the object.

- a) I & II
- b) I, II, and III
- c) I only
- d) II & IV
- e) I & III

A

B

C

D

E

CQ.5.4

A 50 kg box is being slid down a wooden inclined plane with an incline of 55° . If the frictional force it experiences is 80 N, what component of the acceleration parallel to the incline will it achieve? Consider down the plane to be the positive direction.

- a) -6.43 m/s^2
- b) -4.02 m/s^2
- c) 4.02 m/s^2
- d) 6.43 m/s^2

A

B


C

D


E

Activity: Worked Problem

20 . A telephone pole has three cables pulling as shown from above, with $\vec{F}_1 = (300.0\hat{i} + 500.0\hat{j})$, $\vec{F}_2 = -200.0\hat{i}$, and $\vec{F}_3 = -800.0\hat{j}$. (a) Find the net force on the telephone pole in component form. (b) Find the magnitude and direction of this net force.



40 . In the following figure, the horizontal surface on which this block slides is frictionless. If the two forces acting on it each have magnitude $F = 30.0 \text{ N}$ and $M = 10.0 \text{ kg}$, what is the magnitude of the resulting acceleration of the block?



See you next class!

Attribution

This resource was significantly adapted from the Open Stax Instructor Slides provided by Rice University. It is released under a CC-BY 4.0 license.

—— Original resource license ——

OpenStax ancillary resource is © Rice University under a CC-BY 4.0 International license; it may be reproduced or modified but must be attributed to OpenStax, Rice University and any changes must be noted.